

Kent County Influenza Surveillance Report, 2007-2008



CUMULATIVE INFLUENZA CASES OCTOBER 1, 2007 – May 2, 2008 , YEAR TO DATE

Reported Influenza Cases in Kent County, By Type of Influenza, through May 2, 2008

| Type | Frequency | Percent |
|---------|-----------|---------|
| A | 449 | 77.5 % |
| B | 118 | 20.4 % |
| A & B | 2 | .3% |
| Missing | 10 | 1.7% |
| Total | 579 | 100% |

Reported Influenza Cases in Kent County, by Age Group, through May 2, 2008

| Age | Frequency | Percent |
|---------|-----------|---------|
| 0-5 | 91 | 15.7 % |
| 6-17 | 84 | 14.5 % |
| 18-64 | 295 | 50.9% |
| 65+ | 106 | 18.3 % |
| Missing | 3 | .5% |
| Total | 579 | 100% |

Reported Influenza Cases in Kent County, by Sex, Week ending May 2, 2008

| Sex | Frequency | Percent |
|---------|-----------|---------|
| Female | 290 | 50.1% |
| Male | 282 | 48.7% |
| Unknown | 7 | 1.2% |
| Total | 579 | 100% |

Number of Reported Cases of Influenza in Kent County, by Specimen Collection Date, October 1, 2007- May 2, 2008

